



(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 972 520 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

19.01.2000 Bulletin 2000/03

(51) Int. Cl.⁷: **A61K 38/18, A61K 47/26**

(21) Application number: **98901044.2**

(86) International application number:
PCT/JP98/00371

(22) Date of filing: **29.01.1998**

(87) International publication number:
WO 98/33514 (06.08.1998 Gazette 1998/31)

(84) Designated Contracting States:

**AT BE CH DE DK ES FI FR GB GR IE IT LI LU NL
PT SE**

• **ICHIKAWA, Hideki,**
Hoechst Marion Roussel Ltd.
Kawagoe-shi, Saitama 350-11 (JP)

(30) Priority: **30.01.1997 JP 1634997**

(74) Representative:
Vieillefosse, Jean-Claude et al
Hoechst Marion Roussel
Département des Brevets
102, Route de Noisy
93235 Romainville Cédex (FR)

(71) Applicant:

Hoechst Marion Roussel, Ltd.
Tokyo 107-8465 (JP)

(72) Inventors:

• **INAGAKI, Mitsuko**
Iruma-shi, Saitama 358 (JP)

(54) **FREEZE-DRIED COMPOSITION OF BONE MORPHOGENETIC PROTEIN HUMAN MP52**

(57) A stable freeze-dried composition of a bone morphogenetic protein human MP52 wherein coloration and shrinking of MP52 during storage and aggregation at the re-dissolution can be prevented. The composition is obtained by mixing MP52 with mannitol at a weight ratio of 1 : 5 to 1 : 50 followed by freeze-drying.

EP 0 972 520 A1

Description**FIELD OF THE INVENTION**

- 5 [0001] The present invention relates to a lyophilized composition of bone morphogenetic factor human MP52 and a preparation process therefor. More specifically, this invention pertains to a lyophilized composition which contains bone morphogenetic factor human MP52 and mannitol and a preparation process therefor.

BACKGROUND OF THE INVENTION

- 10 [0002] The cDNA of bone morphogenetic factor human MP52 was isolated for the first time in 1994 as an osteogenesis-related factor classified as a TGF- β superfamily (Biochem. Biophys. Res. Comm., Vol. 204, No. 2, 1994). Then, an advanced genetic engineering technology has made it possible to prepare bone morphogenetic factor human MP52 without impairing its bone morphogenetic activity (WO96/33215). Bone morphogenetic factor human MP52 is stored
15 under a lyophilized condition. It is, however, accompanied with a drawback that a volume reduction (shrink) occurs during storage and cohesion of powders occurs at the time of reconstitution.

- [0003] With a view to overcoming the above-described problems, amino acids, saccharides or polyhydric alcohols are used for BMP-2 which is a protein classified as the same TGF- β super-family and has properties closest to bone morphogenetic factor human MP52 (JP-A No. HEI 6-508777). The present inventors, therefore, attempted the application
20 of such additives to bone morphogenetic factor human MP52 but could not overcome the above problems. Described specifically, cohesion at the time of reconstitution was observed even if a neutral or basic amino acid such as alanine, valine or lysine was added to bone morphogenetic factor human MP52 in an amount of 0.5 to 2.5% prior to lyophilization. When a saccharide such as sucrose or dextran was added in an amount of 0.5 to 1%, followed by lyophilization, color development to pale yellowish green and shrink were observed from the lyophilized product. When a polyhydric
25 alcohol such as sorbitol was added in an amount of 0.5 to 1%, followed by lyophilization, bone morphogenetic factor human MP52 was dissolved in the period of lyophilization, which made it impossible to prepare a lyophilized product.

DISCLOSURE OF THE INVENTION

- 30 [0004] The present inventors have proceeded with an extensive investigation with a view to overcoming the above-described problems. As a result, it has been found that when mannitol is added to bone morphogenetic factor human MP52, followed by lyophilization, neither coloring or shrink is observed during the storage of the lyophilized product and cohesion does not occur at the time of reconstitution, leading to the completion of the present invention.

- [0005] The present invention, therefore, provides a lyophilized composition of bone morphogenetic factor human
35 MP52 containing bone morphogenetic factor human MP52 and mannitol. As bone morphogenetic factor human MP52 in the present invention, bone morphogenetic factor human MP52 (which may hereinafter be called "rhMP52") which has been prepared by genetic engineering technology disclosed in WO96/33215 is preferably used. As mannitol, that prescribed as D-mannitol in the Japanese Pharmacopoeia is preferably used. Bone morphogenetic factor human MP52 and mannitol are preferably mixed at a weight ratio of 1 : 5 - 50.

- 40 [0006] The composition according to the present invention can be prepared by lyophilizing an aqueous mixture solution of bone morphogenetic factor human MP52 and mannitol by a conventional method. Described specifically, the composition of the present invention is available by adding a predetermined amount of mannitol to an aqueous solution of purified bone morphogenetic factor human MP52, mixing them, filtering the resulting aqueous mixture solution, filling the filtrate in a sterile vial and carrying out lyophilization.

- 45 [0007] The composition of the present invention is administered to a patient in an amount effective for therapeutic treatment after being dissolved in distilled water for injection or weak acid (about pH 3), for example, a hydrochloric acid solution or citric acid buffer, upon use.

- [0008] The preferred amount of mannitol incorporated in the composition of the present invention was determined by a stability test. The stability test was conducted in accordance with the method described in the instruction for the standard operation procedure prepared based on the Japanese Pharmacopoeia XIII, where properties such as appearance and clarity of solution, electrophoresis and water content at the beginning time of the test and 3 months later and ectopic bone formation after 6 months were observed or measured.

- 50 [0009] As a result, no change in the properties such as appearance and clarity of solution were observed at the beginning time of the test and 3 months later. From the results of the measurement on the electrophoresis and water content, the preparation containing mannitol at the above-described weight ratio was stable both at the beginning time of the test and 3 months later.

- 55 [0010] It is said that the preferred water content of a lyophilized product is generally 2% or lower. It was judged from the above findings that a bone morphogenetic factor human MP52 composition containing mannitol in an amount of 5

to 50 mg, desirably 10 mg, per 1 mg of bone morphogenetic factor human MP52 is preferred as a pharmaceutical product.

[0011] In addition, the ectopic bone formation of each of lyophilized compositions of bone morphogenetic factor human MP52 containing 10, 25 and 50 mg of mannitol, respectively was measured after stored for 6 months. As a result, ectopic bone formation was observed from any composition regardless of the storage temperature or amount of mannitol. Based on the above-described test results, it has been confirmed that the addition of mannitol to bone morphogenetic factor human MP52 prior to lyophilization does not have adverse effects on the bone morphogenetic factor human MP52 and the resulting composition remains stable for a long period of time.

BEST MODE FOR CARRYING OUT THE INVENTION

[0012] The present invention will hereinafter be described more specifically by the following examples.

Example 1 Preparation of an rhMP52 composition

[0013] To 1 mg/ml of an aqueous solution of purified rhMP52, which had been obtained by the preparation process disclosed in WO96/33215. D-mannitol of the Japanese Pharmacopoeia was added in an amount of 10, 25 and 50 mg, respectively, and they were mixed. After the resulting mixture was filtered through a 0.22 μ m membrane filter, 1 ml portions of the filtrate so obtained were filled in vials sterily. They were lyophilized, whereby a composition of the present invention was prepared in the form of pharmaceutical product.

Example 2 Stability test of the rhMP52 composition

[0014] The rhMP52 lyophilized composition obtained in Example 1 was filled in a vial bottle (air-tight, transparent) and was stored at 2 - 8°C, 25°C and 40°C, respectively. The stability was evaluated based on the following criteria after three-month storage. The criteria for evaluation are as follows:

Properties:

[0015]

(Appearance): The composition which remained in the form of a white cake and was not colored was judged as "not changed".

(Clarity of solution): A solution having the composition dissolved in 1 ml of distilled water for injection was judged as "not changed" when it was colorless, transparent and cohesion-free.

(Electrophoresis): A purity of a main band was calculated from an area percentage, after introducing a picture by a film scanner which adopts transmission using a red film and then finding an integration optical density (IOD) % of each band.

(Water content): The water content of the composition was measured by a micro-moisture meter.

[0016] As a result, concerning properties, the composition of the present invention remained unchanged in appearance and clarity of solution, whereas cohesion was observed in a lyophilized product of rhMP52 alone. As a result of electrophoresis, each composition showed good stability (%).

[0017] The measurement results of water content (%) are shown in Table 1 which shows that there is not a large change in the water content (%) among the compositions. Table 1 shows that compared with the product composed only of rhMP52, the water content of each of the compositions according to the present invention is lower at the time of preparation or after storage and therefore shrink does not occur easily.

Table 1

composition	Initial	3 months later		
		2 - 8°C	25°C	40°C
rbMP52 alone	9.7%	9.2%	7.6%	7.8%
+ Mannitol, 10 mg	1.6%	1.6%	1.4%	1.2%
+ Mannitol, 25 mg	0.9%	0.7%	0.6%	0.5%

Table 1 (continued)

composition	Initial	3 months later		
		2 - 8°C	25°C	40°C
+ Mannitol, 50 mg	0.6%	0.5%	0.4%	0.4%

Example 3 Ectopic bone formation of rhMP52

[0018] One vial (1 mg/vial) containing the rhMP52 composition obtained in Example 1 was stored at 4°C and 25°C for 6 months, respectively, then, 1 ml of distilled water for injection was added to the vial, whereby a solution for administration was prepared. The solution thus prepared was intramuscularly administered to an ICR mouse (purchased from Nippon Crea Co., Ltd.) in an amount of 20 µg/20 µl. Two weeks later, the presence or absence of ectopic bone formation was observed through a soft X-ray photographing (n=2). The results are shown in Table 2.

Table 2

composition	4°C	25°C
rhMP52 alone	ebf* was observed	ebf was observed
+ Mannitol, 10 mg	ebf was observed	ebf was observed
+ Mannitol, 25 mg	ebf was observed	ebf was observed
+ Mannitol, 50 mg	ebf was observed	ebf was observed
ebf: ectopic bone formation		

INDUSTRIAL APPLICABILITY

[0019] The lyophilized product of bone morphogenetic factor human MP52 involves problems that it is colored or it shrinks during storage and it coheres at the time of reconstitution. The lyophilized composition according to the present invention, however, is free from such problems. Bone morphogenetic factor human MP52 in the lyophilized composition of the present invention remains stable and no substantial change is observed in purity, water content and ectopic bone formation even after storage for a long time. Thus, the present invention is applicable in the field of pharmaceutical product.

Claims

1. A lyophilized composition of bone morphogenetic factor human MP52, which comprises bone morphogenetic factor human MP52 and mannitol.
2. The lyophilized composition according to claim 1, wherein said bone morphogenetic factor human MP52 and mannitol are mixed at a weight ratio of 1 : 5-50.
3. The lyophilized composition according to claim 1 or 2, wherein said bone morphogenetic factor human MP52 is produced by means of a genetic engineering technology.
4. A process for the preparation of a lyophilized composition of bone morphogenetic factor human MP52, which comprises adding mannitol to an aqueous solution of purified bone morphogenetic factor human MP52 and then lyophilizing the resulting aqueous mixture solution.
5. The process for the preparation of the lyophilized composition according to claim 4, wherein said bone morphogenetic factor human MP52 and mannitol are mixed at a weight ratio of 1 : 5-50.
6. The process for the preparation of the lyophilized composition according to claim 4 or 5, wherein said bone morphogenetic factor human MP52 is produced by means of a genetic engineering technology.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP98/00371

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁶ A61K38/18, A61K47/26 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁶ A61K38/18 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO, 96/33215, A1 (Hoechst Japan Ltd.), October 24, 1996 (24. 10. 96) & AU, 9653470, A	1-6
A	JP, 7-503847, A (Biopharm Gesellschaft zur Biotechnologischen Entwicklung von PharmakambH.), April 27, 1995 (27. 04. 95) & WO, 93/16099, A2 & EP, 625989, A1	1-6
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search April 9, 1998 (09. 04. 98)		Date of mailing of the international search report April 21, 1998 (21. 04. 98)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)